Cornwall-Jackiw-Tomboulis effective potential for quark propagator in real-time thermal field theory and Landau gauge*

Zhou Bang-Rong
College of Physical Sciences, Graduate School of the Chinese Academy of Sciences, Beijing 100049, China and
CCAST (World Laboratory), P.O.Box 8730, Beijing 100080, China

(Dated: March 30, 2005)

We complete the derivation of the Cornwall-Jackiw-Tomboulis effective potential for quark propagator at finite temperature and finite quark chemical potential in the real-time formalism of thermal field theory and in Landau gauge. In the approximation that the function $A(p^2)$ in inverse quark propagator is replaced by unity, by means of the running gauge coupling and the quark mass function invariant under the renormalization group in zero temperature Quantum Chromadynamics (QCD), we obtain a calculable expression for the thermal effective potential which will be a useful means to research chiral phase transition in QCD in the real-time formalism.

PACS numbers: 11.10.Wx; 12.38.Lg; 11.30.Rd; 12.38.Aw
Keywords: CJT effective potential, real-time thermal field theory, QCD, quark propagator, Landau gauge

arXiv:hep-th/0512044v1 5 Dec 2005

Effective potential [1, 2, 3, 4, 5] is a basic means to research the vacuum of quantum field theory and spontaneous symmetry breaking. For discussion of chiral symmetry phase transition in QCD, one must generalize the Cornwall-Jackiw-Tomboulis (CJT) effective potential [6] for quark propagator to finite temperature and finite quark chemical potential case. At present, some work on the thermal CJT effective potential are mainly based on the imaginary-time formalism [7], very few were found in literature which are based on the real-time formalism [8]. In fact, in the real-time formalism, besides that it is no longer necessary to do analytic continuation of the discrete Matsubara frequency, one can directly use the well-known results in zero temperature QCD, e.g. the running gauge coupling and the running quark mass function consistent with the renormalization group (RG) analysis before thermalizing propagators by thermal transformation matrices, and this will certainly bring great advantages for practical calculation.

In a preceding paper [9], for deriving the thermal Schwinger-Dyson (SD) equation for quark self-energy, we gave a formal expression for the thermal CJT effective potential for quark propagator in Landau gauge. In this paper, we will continue to reduce the formal expression to an explicit formula for future use.

For convenience of the derivation and self-containment of the paper, we will first put down the relevant results given in Ref. [9]. The formal expression for the CJT effective potential for quark propagator at finite temperature T and finite quark chemical potential μ has been expressed by

$$V[G,G^*]=V_1[G,G^*]+V_2[G,G^*] \tag{1}$$

where in momentum space the contribution from one-loop diagrams

$$\begin{aligned}V_1[G,G^*] \;=\; & iN_fN_c\Big(\big\langle \cos^2\theta_p\big\{\mathrm{tr}\ln\big[S(p)G^{-1}(p)\big]+\mathrm{tr}\big[S^{-1}(p)G(p)\big]\big\}\big\rangle \\ & +\big\langle \sin^2\theta_p\big\{\mathrm{tr}\ln\big[S^*(p)G^{*-1}(p)\big]+\mathrm{tr}\big[S^{*-1}(p)G^*(p)\big]\big\}\big\rangle-\big\langle \mathrm{tr}1\big\rangle\Big),\end{aligned} \tag{2}$$

and the contribution from two-loop vacuum diagrams

$$\begin{aligned}V_2[G,G^*] \;=\; & \frac{i}{2}N_fN_cC_2(\underline{N}_c)\big\langle\big\langle g^2\big[(p-q)^2\big]\mathrm{tr}\big[\gamma^\mu G_T^{11}(p)\gamma^\nu G_T^{11}(q)\big]\big[D_{\mu\nu}(p-q)\big]_T^{11}\big\rangle\big\rangle \\ & -\frac{i}{2}N_fN_cC_2(\underline{N}_c)\big\langle\big\langle g^2\big[(p-q)^2\big]\mathrm{tr}\big[\gamma^\mu G_T^{12}(p)\gamma^\nu G_T^{21}(q)\big]\big[D_{\mu\nu}(p-q)\big]_T^{12}\big\rangle\big\rangle.\end{aligned} \tag{3}$$

In the above denotations, $\langle\cdots\rangle$ and $\langle\langle\cdots\rangle\rangle$ represent respectively the integration $\int d^4p/(2\pi)^4$ and $\int d^4pd^4q/(2\pi)^8$, N_f and N_c are respectively the number of flavor and color of the quarks, $C_2(\underline{N}_c)=(N_c^2-1)/2N_c$,

$$S(p)=i/(\not p+i\varepsilon)\ \text{ and }\ S^*(p)=-i/(\not p-i\varepsilon)\ \text{ with }\ \not p=\gamma^\mu p_\mu \tag{4}$$

---

* The project supported by the National Natural Science Foundation of China.

are the propagator of massless quark and its conjugate,

$$G(p)=i/\big[\,\not p-m(p^2)+i\varepsilon\big]\ \text{ and }\ G^*(p)=-i/\big[\,\not p-m(p^2)-i\varepsilon\big] \tag{5}$$

are complete quark propagator and its conjugate, in which we have made the approximation $A(p^2)=1$ on the basis of the arguments given in Sect.IV of Ref. [9] and the replacement $B(p^2)\to m(p^2)$, $g^2\big[(p-q)^2\big]$ is the running gauge coupling, tr means the trace of spinor matrices,

$$\sin^2\theta_p=\theta(p^0)\tilde{n}(p^0-\mu)+\theta(-p^0)\tilde{n}(-p^0+\mu)\ \text{ with }\ \tilde{n}(p^0-\mu)=1/\big[e^{\beta(p^0-\mu)}+1\big]\ \text{ and }\ \beta=1/T, \tag{6}$$

$$G_T^{11}(p)=\cos^2\theta_pG(p)-\sin^2\theta_pG^*(p),\ \ G_T^{12}(p)=-\cos\theta_p\ \sin\theta_p\ e^{\beta\mu/2}\big[G(p)+G^*(p)\big]=-e^{\beta\mu}G_T^{21}(p), \tag{7}$$

and

$$\big[D_{\mu\nu}(p-q)\big]_T^{11}=D_{\mu\nu}(p-q)-g_{\mu\nu}2\pi n(p^0-q^0)\delta\big[(p-q)^2\big],\ \ \big[D_{\mu\nu}(p-q)\big]_T^{12}=-g_{\mu\nu}e^{\beta|p^0-q^0|/2}2\pi n(p^0-q^0)\delta\big[(p-q)^2\big] \tag{8}$$

with

$$D_{\mu\nu}(p-q)=\frac{-i}{(p-q)^2+i\varepsilon}\Delta_{\mu\nu}(p-q),\ \ \Delta_{\mu\nu}(p-q)=g_{\mu\nu}-\frac{(p-q)_\mu(p-q)_\nu}{(p-q)^2+i\varepsilon} \tag{9}$$

and

$$n(p^0-q^0)=1/\big[e^{\beta|p^0-q^0|}-1\big]. \tag{10}$$

Now we begin performing detailed calculations of $V[G,G^*]$. First rewrite $V_1[G,G^*]$ in Eq.(2) by

$$V_1[G,G^*]=V_{10}+V_{1T} \tag{11}$$

with

$$V_{10}=iN_fN_c\Big(\Big\langle\mathrm{tr}\ln\big[S(p)G^{-1}(p)\big]+\mathrm{tr}\big[S^{-1}(p)G(p)\big]-\mathrm{tr}1\Big\rangle\Big) \tag{12}$$

and

$$V_{1T}=iN_fN_c\Big\langle\sin^2\theta_p\big\{\mathrm{tr}\ln\big[S^*(p)G^{*-1}(p)\big]+\mathrm{tr}\big[S^{*-1}(p)G^*(p)\big]-\mathrm{tr}\ln\big[S(p)G^{-1}(p)\big]-\mathrm{tr}\big[S^{-1}(p)G(p)\big]\big\}\Big\rangle. \tag{13}$$

By means of Eqs.(4) and (5) we get

$$\begin{aligned}
V_{10}&=iN_fN_c\Big\langle\mathrm{tr}\left\{\ln\left[1-\frac{m(p^2)}{\not p+i\varepsilon}\right]+\frac{m(p^2)}{\not p-m(p^2)+i\varepsilon}\right\}\Big\rangle\\
&=iN_fN_c\Big\langle\mathrm{tr}\left\{\sum_{k=1}^{\infty}\frac{-1}{k}\left[\frac{m(p^2)}{\not p+i\varepsilon}\right]^k+\frac{m(p^2)}{\not p-m(p^2)+i\varepsilon}\right\}\Big\rangle\\
&=i2N_fN_c\int\frac{d^4p}{(2\pi)^4}\left\{\ln\left[1-\frac{m^2(p^2)}{p^2+i\varepsilon}\right]+\frac{2m^2(p^2)}{p^2-m^2(p^2)+i\varepsilon}\right\}\\
&=-2N_fN_c\int\frac{d^4\bar{p}}{(2\pi)^4}\left\{\ln\frac{\bar{p}^2+m^2(\bar{p}^2)}{\bar{p}^2}-\frac{2m^2(\bar{p}^2)}{\bar{p}^2+m^2(\bar{p}^2)}\right\}
\end{aligned} \tag{14}$$

In the last line we have made the Wick rotation and the changes of variables

$$\bar{p}^0=ip^0,\ \bar{p}^i=p^i\ (i=1,2,3). \tag{15}$$

In similar way we obtain

$$\begin{aligned}
V_{1T}&=i2N_fN_c\Big\langle\sin^2\theta_p\left\{\ln\left[1+\frac{m^2(p^2)}{p^2-m^2(p^2)+i\varepsilon}\right]-\ln\left[1+\frac{m^2(p^2)}{p^2-m^2(p^2)-i\varepsilon}\right]\right.\\
&\qquad\left.-2p^2\left[\frac{1}{p^2-m^2(p^2)+i\varepsilon}-\frac{1}{p^2-m^2(p^2)-i\varepsilon}\right]\right\}\Big\rangle\\
&=-4N_fN_c\Big\langle\sin^2\theta_p\mathrm{Im}\left\{\ln\left[1+\frac{m^2(p^2)}{p^2-m^2(p^2)+i\varepsilon}\right]-\frac{2p^2}{p^2-m^2(p^2)+i\varepsilon}\right\}\Big\rangle\\
&=-4N_fN_c\Big\langle\sin^2\theta_p\Big\{-\pi m^2(p^2)\delta\big[p^2-m^2(p^2)\big]\sum_{k=1}^{\infty}\left(\frac{-m^2(p^2)\big[p^2-m^2(p^2)\big]}{\big[p^2-m^2(p^2)\big]^2+\varepsilon^2}\right)^{k-1}+2\pi p^2\delta\big[p^2-m^2(p^2)\big]\Big\}\Big\rangle\\
&=-4\pi N_fN_c\Big\langle\sin^2\theta_pm^2(p^2)\delta\big[p^2-m^2(p^2)\big]\Big\rangle,
\end{aligned} \tag{16}$$

where the relation $\delta\big[p^2-m^2(p^2)\big]\big[p^2-m^2(p^2)\big]=0$ has been used.

Next we rewrite $V_2[G,G^*]$ in Eq.(3) in terms of Eqs.(7)-(9) by

$$V_2[G,G^*]=V_{20}+V_{2T}, \tag{17}$$

where V_{20} and V_{2T} are respectively the contributions from the two-loop vacuum diagrams without and with thermal corrections of propagators. V_{20} can be expressed by

$$\begin{aligned} V_{20} &= -\frac{1}{2}N_fN_cC_2(\underline{N}_c)\Big\langle\Big\langle g^2\big[(p-q)^2\big]\mathrm{tr}\left\{\gamma^\mu\big[\,\not p+m(p^2)\big]\gamma^\nu\big[\,\not q+m(q^2)\big]\right\}\frac{1}{p^2-m^2(p^2)+i\varepsilon}\cdot\frac{1}{q^2-m^2(q^2)+i\varepsilon} \\ &\quad\cdot\frac{1}{(p-q)^2+i\varepsilon}\Delta_{\mu\nu}(p-q)\Big\rangle\Big\rangle \\ &= -2N_fN_cC_2(\underline{N}_c)\Big\langle\Big\langle g^2\big[(p-q)^2\big]\big[3m(p^2)m(q^2)+(p-q)^2E(p,q)\big]\frac{1}{p^2-m^2(p^2)+i\varepsilon}\cdot\frac{1}{q^2-m^2(q^2)+i\varepsilon} \\ &\quad\cdot\frac{1}{(p-q)^2+i\varepsilon}\Big\rangle\Big\rangle \\ &= -6N_fN_cC_2(\underline{N}_c)\int\frac{d^4\bar p d^4\bar q}{(2\pi)^8}g^2\big[\max(\bar p^2,\bar q^2)\big]\frac{m(\bar p^2)}{\bar p^2+m^2(\bar p^2)}\cdot\frac{m(\bar q^2)}{\bar q^2+m^2(\bar q^2)}\cdot\frac{1}{(\bar p-\bar q)^2} \end{aligned}\tag{18}$$

where we have used the result

$$\mathrm{tr}\Big\{\gamma^\mu\big[\,\not p+m(p^2)\big]\gamma^\nu\big[\,\not q+m(q^2)\big]\Big\}\Delta_{\mu\nu}(p-q)=4\big[3m(p^2)m(q^2)+(p-q)^2E(p,q)\big] \tag{19}$$

with the denotation

$$E(p,q)=1-\frac{1}{2}\left[p^2+q^2+\frac{(p^2-q^2)^2}{(p-q)^2}\right]/(p-q)^2, \tag{20}$$

and after the Wick rotation, the running gauge coupling $g^2\big[(\bar p-\bar q)^2\big]$ has been approximated by [10]

$$g^2\big[\max(\bar p^2,\bar q^2)\big]=\theta(\bar p-\bar q)g^2(\bar p^2)+\theta(\bar q-\bar p)g^2(\bar q^2) \tag{21}$$

so that

$$\int d\Omega_{\bar q}E(\bar p,\bar q)=0. \tag{22}$$

By using Eq.(19) and the result that

$$\mathrm{tr}\Big\{\gamma^\mu\big[\,\not p+m(p^2)\big]\gamma^\nu\big[\,\not q+m(q^2)\big]\Big\}g_{\mu\nu}=4\,\big[4m(p^2)m(q^2)-2p\cdot q\big] \tag{23}$$

and considering some symmetry of the integrand under the exchange between the variables p and q we may express V_{2T} by

$$\begin{aligned} V_{2T} &= i4\pi N_fN_cC_2(\underline{N}_c)g^2(0)\Big\langle\Big\langle\big[4m(p^2)m(q^2)-p^2-q^2\big]n(p^0-q^0)\delta\big[(p-q)^2\big] \\ &\quad\left\{\frac{1}{p^2-m^2(p^2)+i\varepsilon}\cdot\frac{1}{q^2-m^2(q^2)+i\varepsilon}+i4\pi\sin^2\theta_p\delta\big[p^2-m^2(p^2)\big]\mathrm{P}\frac{1}{q^2-m^2(q^2)}\right\}\Big\rangle\Big\rangle \\ &\quad-8\pi N_fN_cC_2(\underline{N}_c)\Big\langle\Big\langle\frac{g^2\big[(p-q)^2\big]}{(p-q)^2+i\varepsilon}\big[3m(p^2)m(q^2)+(p-q)^2E(p,q)\big]\sin^2\theta_p\delta\big[p^2-m^2(p^2)\big] \\ &\quad\left\{\frac{i}{q^2-m^2(q^2)+i\varepsilon}-\pi\sin^2\theta_q\delta\big[q^2-m^2(q^2)\big]\right\}\Big\rangle\Big\rangle+\Delta V_{2T}, \end{aligned}\tag{24}$$

where "P" in the second line of Eq.(24) means the principle value of the integration and

$$\begin{aligned} \Delta V_{2T} &= -i16\pi^3N_fN_cC_2(\underline{N}_c)\Big\langle\Big\langle g^2\big[(p-q)^2\big]\big[4m(p^2)m(q^2)-p^2-q^2\big]\delta\big[p^2-m^2(p^2)\big]\delta\big[q^2-m^2(q^2)\big]\delta\big[(p-q)^2\big] \\ &\quad n(p^0-q^0)\left\{\sin^2\theta_p\sin^2\theta_q-\sin^2\theta_p+\sin\theta_p\cos\theta_p\sin\theta_q\cos\theta_qe^{\beta|p^0-q^0|/2}\right\}\Big\rangle\Big\rangle \end{aligned}\tag{25}$$

Let $p^2=m_1^2$ is a real root of the equation $p^2=m^2(p^2)$, i.e. m_1 obeys the equation $m_1^2=m^2(m_1^2)$, then we will have $\delta\big[p^2-m^2(p^2)\big]=\delta(p^2-m_1^2)/f(m_1^2)$ with $f(m_1^2)=|1-\partial m^2(p^2)/\partial p^2|_{p^2=m_1^2}$. Thus we get

$$\begin{aligned}\Delta V_{2T} &= -i32\pi^3 N_f N_c C_2(\underline{N}_c) g^2(0)\frac{m_1^2}{f^2(m_1^2)}\Big\langle\Big\langle \delta(p^2-m_1^2)\delta(q^2-m_1^2)\delta\big[(p-q)^2\big] \\ &\qquad n(p^0-q^0)\Big\{\sin^2\theta_p\sin^2\theta_q-\sin^2\theta_p+\sin\theta_p\cos\theta_p\sin\theta_q\cos\theta_q e^{\beta|p^0-q^0|/2}\Big\}\Big\rangle\Big\rangle \\ &= -i32\pi^3 N_f N_c C_2(\underline{N}_c) g^2(0)\frac{m_1^2}{f^2(m_1^2)}\Big\langle\Big\langle \delta\big[(p+q)^2-m_1^2\big]\delta(q^2-m_1^2)\delta(p^2) \\ &\qquad n(p^0)\Big\{\sin^2\theta_{p+q}\sin^2\theta_q-\sin^2\theta_{p+q}+\sin\theta_{p+q}\cos\theta_{p+q}\sin\theta_q\cos\theta_q e^{\beta|p^0|/2}\Big\}\Big\rangle\Big\rangle \qquad (26)\end{aligned}$$

In the second equality above, we have made the change of variables $p-q\rightarrow p$. It is indicated that the condition in which the three δ functions in Eq.(26) are not equal to zeroes simultaneously is $p^0=|\vec{p}|=0$. Taking this and the definition of $\sin^2\theta_p$ given by Eq.(6) into account, we are led to that

$$\big[\sin^2\theta_{p+q}\sin^2\theta_q-\sin^2\theta_{p+q}+\sin\theta_{p+q}\cos\theta_{p+q}\sin\theta_q\cos\theta_q e^{\beta|p^0|/2}\big]\big|_{p^0=0}=\sin^2\theta_q\big[\sin^2\theta_q-1+\cos^2\theta_q\big]=0.$$

On the other hand, it is also noted that because

$$\int d^4p\,\delta(p^2)n(p^0)=\int dp^0\int d\Omega_{\vec{p}}\int d|\vec{p}||\vec{p}|^2\frac{1}{2|\vec{p}|}\big[\delta(p^0-|\vec{p}|)+\delta(p^0+|\vec{p}|)\big]\frac{1}{e^{\beta|\vec{p}|}-1},$$

no singularity could appear in integrand in Eq.(26) when $|\vec{p}|\rightarrow 0$. Hence we can conclude that

$$\Delta V_{2T}=0. \qquad (27)$$

In summary, from Eqs.(1),(11),(14),(16),(17),(18) (24) and (27) we may write the final expression for the CJT effective potential for quark propagator in the approximation $A(p^2)=1$ at finite temperature and finite chemical potential in the real-time formalism and in Landau gauge as follows.

$$V[G,G^*]=V_0+V_T \qquad (28)$$

$$\begin{aligned}V_0 &= V_{10}+V_{20} \\ &= -2N_f N_c\Big\langle \ln\frac{\bar{p}^2+m^2(\bar{p}^2)}{\bar{p}^2}-\frac{2m^2(\bar{p}^2)}{\bar{p}^2+m^2(\bar{p}^2)}\Big\rangle_E \\ &\quad -6N_f N_c C_2(\underline{N}_c)\Big\langle\Big\langle g^2[\max(\bar{p}^2,\bar{q}^2)]\frac{m(\bar{p}^2)}{\bar{p}^2+m^2(\bar{p}^2)}\cdot\frac{m(\bar{q}^2)}{\bar{q}^2+m^2(\bar{q}^2)}\cdot\frac{1}{(\bar{p}-\bar{q})^2}\Big\rangle\Big\rangle_E, \qquad (29)\end{aligned}$$

where $\langle\cdots\rangle_E$ and $\langle\langle\cdots\rangle\rangle_E$ means the integrations of the Euclidean momentums $\int d^4\bar{p}/(2\pi)^4$ and $\int d^4\bar{p}d^4\bar{q}/(2\pi)^8$ respectively.

$$\begin{aligned}V_T &= V_{1T}+V_{2T} \\ &= -4\pi N_f N_c\Big\langle \sin^2\theta_p m^2(p^2)\delta\big[p^2-m^2(p^2)\big]\Big\rangle \\ &\quad +i4\pi N_f N_c C_2(\underline{N}_c)g^2(0)\Big\langle\Big\langle\big[4m(p^2)m(q^2)-p^2-q^2\big]n(p^0-q^0)\delta\big[(p-q)^2\big] \\ &\quad \Big\{\frac{1}{p^2-m^2(p^2)+i\varepsilon}\cdot\frac{1}{q^2-m^2(q^2)+i\varepsilon}+i4\pi\sin^2\theta_p\delta\big[p^2-m^2(p^2)\big]\mathrm{P}\frac{1}{q^2-m^2(q^2)}\Big\}\Big\rangle\Big\rangle \\ &\quad -8\pi N_f N_c C_2(\underline{N}_c)\Big\langle\Big\langle\frac{g^2\big[(p-q)^2\big]}{(p-q)^2+i\varepsilon}\big[3m(p^2)m(q^2)+(p-q)^2E(p,q)\big]\sin^2\theta_p\delta\big[p^2-m^2(p^2)\big] \\ &\quad \Big\{\frac{i}{q^2-m^2(q^2)+i\varepsilon}-\pi\sin^2\theta_q\delta\big[q^2-m^2(q^2)\big]\Big\}\Big\rangle\Big\rangle \qquad (30)\end{aligned}$$

In Eqs.(29) and (30), the running gauge coupling constant $g^2\big[(p-q)^2\big]$ and the running quark mass function $m(p^2)$ can be taken from the known results in zero temperature QCD which are consistent with RG analysis. After introducing the infrared momentum cutoff p_c^2, we can write [11]

$$g^2\big[(p-q)^2\big]=2\pi^2 A/\ln\frac{|(p-q)^2|+p_c^2}{\Lambda_{QCD}^2} \qquad (31)$$

and

$$m(p^2) = \hat{m}\left(\ln\frac{|p^2|+p_c^2}{\Lambda_{QCD}^2}\right)^{-A/2} + \frac{2\pi^2 A}{3}\frac{\phi}{(|p^2|+p_c^2)\varepsilon(p^2)}\left(\ln\frac{|p^2|+p_c^2}{\Lambda_{QCD}^2}\right)^{A/2-1}, \quad \varepsilon(p^2) \equiv \frac{p^2}{|p^2|}, \tag{32}$$

where $A = 24/(33-2N_f)$, Λ_{QCD} is the RG-invariant mass scale parameter, $\phi \equiv \langle 0|\bar{\psi}\psi|0\rangle/(\ln\mu^2/\Lambda_{QCD}^2)^{A/2}$ and $\hat{m} = m[\ln(\mu^2/\Lambda_{QCD}^2)]^{A/2}$ are respectively the RG-invariant quark-antiquark condensates and the current quark mass, and μ is the renormalization scale parameter. Eqs. (28)-(30) combined with Eqs. (31)-(32) give a calculable expression for the thermal CJT effective potential for quark mass function $m(p^2)$. Obviously, it is a function of temperature T, quark chemical potential μ, the quark-antiquark condensates ϕ and, of course, the current quark mass $\hat{m}$ as well. The result provides a useful means to research chiral symmetry phase transition in QCD in the real-time formalism of thermal field theory.

---

[1] W. Heisenberg and H. Euler, Z. Phys. **98** (1936) 714; J. Schwinger, Phys. Rev. **82** (1951) 664.

[2] J. Goldstone, A. Salam, and J. Weinberg, Phys. Rev. **127** (1962) 965; G. Jona-Lasinio, N. Cimento **34** (1964) 1790.

[3] S. Coleman and E. Weinberg, Phys. Rev. D**7** (1973) 1888.

[4] R. Jackiw, Phys. Rev. D**9** (1974) 1686.

[5] J. Iliopoulos, C. Itzykson, and A. Martin, Rev. Mod. Phys. **47** (1975) 165.

[6] J. M. Cornwall, R. Jackiw, and E. Tomboulis, Phys. Rev. D**10** (1974) 2428.

[7] O. Kiriyama, M. Maruyama, and F. Takagi, Phys. Rev. D**62** (2000) 105008.

[8] N. P. Landsman and Ch. G. van Weert, Phys. Rep. **145** (1987) 141.

[9] Zhou Bang-Rong, Real-time thermal Schwinger-Dyson equation for quark self-energy in Landau gauge, Commun. Theor. Phys. to appear.

[10] K.Higashijima, Phys. Lett. **124B** (1983)257; V.A.Miransky, Yad. Fiz. **38** (1983) 468 [Sov.J.Nucl. Phys. 38 (1983) 280].

[11] K. Higashijima, Prog. Theor. Phys. Suppl. **104** (1991) 1.